



Apr 16th, 9:00 AM - 10:00 AM

# Determining the Viability of Preserved Cell Suspensions Frozen Over Time with Varied Glycerol Concentrations

Elyse McCormick

*Illinois Wesleyan University*

Brian Walter, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

 Part of the [Biology Commons](#), and the [Education Commons](#)

Elyse McCormick and Brian Walter, Faculty Advisor, "Determining the Viability of Preserved Cell Suspensions Frozen Over Time with Varied Glycerol Concentrations" (April 16, 2016). *John Wesley Powell Student Research Conference*. Paper 11.

<http://digitalcommons.iwu.edu/jwprc/2016/posters/11>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P25

**DETERMINING THE VIABILITY OF PRESERVED CELL  
SUSPENSIONS FROZEN OVER TIME WITH VARIED  
GLYCEROL CONCENTRATIONS**

Elyse McCormick and Brian Walter\*  
Biology Department, Illinois Wesleyan University

Cell Viability measures how long cells will stay alive after being cultured, that is, how long they are viable in a given experimental condition. In this project, cells were subjected to a -70° C freezer and different concentrations of glycerol as a cryoprotectant. Two organisms were tested in this experiment: *Schizochytrium* spp. and *E. coli*. Vial lots of each organism were made and placed in a -70°C freezer. After allotted times, vials with concentrations of 10% glycerol, 15% glycerol, and 20% glycerol added to the broth were tested by aerobic plate counts to see the viability of the cells. The *Schizochytrium* spp. vials were tested by a Cellometer® cell counter as well. The preliminary results seem to show that this experiment is a good and useful model for studying viability in these strains. Ultimately, this experiment will be run for two years in order to determine viability over time.